


Testing Variables for Monitoring Estuarine Nutrient Enrichment within North Atlantic Parks: Measuring Agents of Change

*Luke Cole, Scott Nixon, Steve Granger
Graduate School of Oceanography, URI*

Coordinator: Bryan Milstead, NPS



Nitrogen Inputs to Marine Systems




Acadia National Park, ME
<http://rosswarner.com/131b.jpg>

Parks Involved

- **Acadia National Park, ME**
 - ACAD
- **Cape Cod National Seashore, MA**
 - CACO
- **Fire Island National Seashore, NY**
 - FIIS
- **Gateway National Recreation Area, NY/NJ**
 - GATE
- **Assateague Island National Seashore, MD**
 - ASIS
- **Colonial National Historical Park, VA**
 - COLO


Parks In NPS Study



Acadia National Park, ME

Cape Cod National Seashore, MA

Parks In NPS Study





Fire Is. National Seashore, NY



Gateway Natl. Rec. Area, NY/NJ

Parks In NPS Study



Assateague Is. Natl. Seashore, MD



Colonial Natl. Historical Park, VA


Project Goals

- **Delineate a Sphere of Influence**
- **Establish a GIS Inventory for each Park**
 - Watershed/Elevation
 - Land Use
 - Census
 - Fertilizer/Manure
 - Livestock
 - Soils
- **Assemble Pertinent Non-GIS Data**
 - Fertilizer use
 - Livestock coverage
 - Crop Coverage
 - Water consumption
 - WWTP discharges
 - Direct Atmospheric Deposition
- **Apply Nutrient Models (NLM, MANAGE)**
- **Analyze Historical Data**

Spheres of Influence

A sphere of influence was delineated around each park based upon:


- Watershed
- Coastline
- Political Boundaries
- Water Bodies



GIS Content

Land Use and Elevation

Example: *Acadia National Park*



Land Use

+



Elevation Model



=



GIS Content

Census Data

Example: *Fire Island National Seashore*



Note: Data from
2000 US Census and
is normalized by area
(people ha⁻¹)

Modeling

- **Assemble GIS and numerical data**
- **Models to be used:**

- **Valiela N-Loading Model**

Valiela, I., G. Collins, J. Kremer, K. Lajtha, M. Geist, B. Seely, J. Brawley, and C. H. Sham. 1997. Nitrogen loading from coastal watersheds to receiving estuaries: new method and application. *Ecological Applications* 7(2):358-380.

- **MANAGE N-Source Model**

Kellogg, D.Q., L. Joubert, and A.J. Gold. 1997. MANAGE: A Method for Assessment, Nutrient-loading, And Geographic Evaluation of Nonpoint Pollution. University of Rhode Island Cooperative Extension, Department of Natural Resources Science. Kingston, RI.

The Models

- **Valiela N-Loading Model (NLM)**
 - Original Use: Waquoit Bay, Cape Cod, MA
 - Assumptions: Similar soil type
Minimal Agriculture
 - Method: Area-weighted values (e.g. land use)
 - Calculation: Excel Spreadsheet
 - Output: Total N-loading (kg N y^{-1})

The Models

- **MANAGE N-Loading**


- Original Use: Southern NE (MA and RI)
- Assumptions: No decay or retention
- Method: Area-weighted values (e.g. land use)

Loading Coefficients


- Calculation: GIS and Excel Spreadsheet
- Output: Total N-source (kg N y^{-1})

Historical Analysis

- Data compilation
 - GIRAS Land Use
- GIS Analysis
- Model Analysis



FIIS - Historic Land Use - GIRAS



FIIS - Present Land Use - LULC

Upcoming Steps

- **Run NLM and MANAGE for each park**
- **Compare and QA/QC Results**
 - NLM: N input with retention
 - MANAGE: Gross N source (no retention)
- **Compile Metadata**
- **Prepare Report**